



CIGNA MEDICAL COVERAGE POLICY

The following Coverage Policy applies to all plans administered by CIGNA Companies including plans administered by Great-West Healthcare, which is now a part of CIGNA.

Subject Abdominoplasty and Panniculectomy

Effective Date 2/15/2009
Next Review Date 2/15/2010
Coverage Policy Number 0027

Table of Contents

Coverage Policy	1
General Background	2
Coding/Billing Information	3
References	4
Policy History	8

Hyperlink to Related Coverage Policies

Bariatric Surgery
Redundant Skin Surgery

INSTRUCTIONS FOR USE

Coverage Policies are intended to provide guidance in interpreting certain **standard** CIGNA HealthCare benefit plans as well as benefit plans formerly administered by Great-West Healthcare. Please note, the terms of a participant's particular benefit plan document [Group Service Agreement (GSA), Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a participant's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a participant's benefit plan document **always supercedes** the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable group benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. Proprietary information of CIGNA. Copyright ©2009 CIGNA

Coverage Policy

Abdominoplasty and panniculectomy are specifically excluded under many benefit plans. In addition, coverage for these services may be subject to the provisions of a cosmetic and/or reconstructive surgery benefit. These procedures are considered cosmetic when performed solely to improve physical appearance. Cosmetic surgery is specifically excluded under many benefit plans. Please refer to the applicable benefit plan document to determine benefit availability and the terms and conditions of coverage.

If coverage for the service is available, the following conditions of coverage.

CIGNA covers panniculectomy, with or without abdominoplasty, as medically necessary when BOTH of the following conditions are met as demonstrated on preoperative photographs:

- Pannus hangs at or below the level of the symphysis pubis
- Pannus causes a chronic and persistent skin condition (e.g., intertriginous dermatitis, panniculitis, cellulitis or skin ulcerations) that is refractory to at least six months of medical treatment. In addition to good hygiene practices, treatment should include topical antifungals, topical and/or systemic corticosteroids, and/or local or systemic antibiotics.

Note: If the procedure is being performed following significant weight loss, in addition to meeting the criteria noted above, there should be evidence that the individual has maintained a stable weight for at

least six months. If the weight loss is the result of bariatric surgery, abdominoplasty/panniculectomy should not be performed until at least 18 months after bariatric surgery and only when weight has been stable for at least the most recent six months.

CIGNA does not cover abdominoplasty or panniculectomy when performed primarily for ANY of the following indications because it is considered not medically necessary (this list may not be all-inclusive):

- treatment of neck or back pain
- improving appearance (i.e., cosmesis)
- repairing abdominal wall laxity or diastasis recti
- treating psychological symptomatology or psychosocial complaints
- when performed in conjunction with abdominal or gynecological procedures (e.g., abdominal hernia repair, hysterectomy, obesity surgery) unless criteria for panniculectomy and abdominoplasty are met separately

CIGNA does not cover suction-assisted lipectomy when performed alone and not as part of a medically necessary and covered abdominoplasty/panniculectomy procedure.

General Background

Abdominoplasty, often referred to as a "tummy tuck," is a surgical procedure that tightens lax anterior abdominal wall muscles and removes excess abdominal skin and fat. This recontouring of the abdominal wall area is often performed solely to improve the appearance of a protuberant abdomen by creating a flatter, firmer abdomen. The standard abdominoplasty involves plication of the anterior rectus sheath for muscle diastasis (i.e., repair of diastasis recti) and removal of excess fat and skin. Traditional abdominoplasty can be performed as an open procedure or endoscopically. Abdominoplasty completed by endoscopic guidance is usually reserved for those patients who seek less extensive contouring of the abdominal wall. Mini-abdominoplasty, with or without liposuction, is a partial abdominoplasty involving the incision of the lower abdomen only. The procedure is generally performed solely for cosmetic purposes in order to improve the appearance of the abdominal area.

Panniculectomy, a procedure closely related to abdominoplasty, is the surgical excision of a redundant, large and/or long overhanging apron of skin and subcutaneous fat located in the lower abdominal area. The condition may accompany significant overstretching of the lax anterior abdominal wall and, hence, often occurs in morbidly obese individuals or following substantial weight loss. The severity of abdominal deformities is graded as follows (American Society of Plastic Surgeons [ASPS], 2007d):

- Grade 1: panniculus covers hairline and mons pubis but not the genitals
- Grade 2: panniculus covers genitals and upper thigh crease
- Grade 3: panniculus covers upper thigh
- Grade 4: panniculus covers mid-thigh
- Grade 5: panniculus covers knees and below

Treatment of this redundant skin and fat is often performed solely for cosmesis, to improve the appearance of the abdominal area. The presence of a massive overhanging apron of fat and skin, however, may result in chronic and persistent local skin conditions in the abdominal folds. These conditions may include intertrigo, intertriginous dermatitis, cellulitis, ulcerations or tissue necrosis, or they may lead to painful inflammation of the subcutaneous adipose tissue (i.e., panniculitis). When panniculitis is severe, it may interfere with activities of daily living, such as personal hygiene and ambulation. In addition to excellent personal hygiene practices, treatment of these skin conditions generally involves topical or systemic corticosteroids, topical antifungals, and topical or systemic antibiotics. Concurrent abdominal and pelvic surgical procedures (e.g., hernia repair, hysterectomy, obesity surgery) may also be performed in the same operative setting as abdominoplasty or panniculectomy.

Diastasis Recti

Abdominoplasty is frequently performed to treat diastasis recti, a condition that involves the separation of the two sides of the rectus abdominis muscles in the midline at the linea alba. Other than its untoward cosmetic appearance, diastasis recti does not lead to any complications that require intervention (Condon, 1995). Diastasis recti has no clinical significance, does not require treatment and is not considered a true hernia. When performed for the sole purpose of repairing diastasis recti, abdominoplasty is considered cosmetic in nature and not medically necessary.

Abdominoplasty and Panniculectomy at the Time of Abdominal or Gynecological Surgeries

Abdominal surgeries such as hernia repair (i.e., incisional/ventral, epigastric or umbilical) or obesity surgery may be performed alone or in combination with abdominoplasty and panniculectomy. In addition, some surgeons perform these procedures at the same time as gynecological or pelvic procedures, such as hysterectomy. Although it has been proposed that performing abdominoplasty or panniculectomy in the same operative session as abdominal or gynecological surgeries may facilitate surgical access or promote postoperative wound healing and minimize the potential for wound complications, such as dehiscence or necrosis, there is insufficient evidence in the published, peer-reviewed scientific literature to support such assertions. Performing an abdominoplasty at the same operative session as abdominal operations (e.g., hernia repair, gastric bypass) or gynecological procedures is not essential for the successful clinical outcome of the abdominal or gynecological surgical procedure. In the absence of chronic and persistent skin conditions or interference with activities of daily living, abdominoplasty and panniculectomy are considered not medically necessary when performed in conjunction with abdominal or pelvic/gynecological surgeries to facilitate surgical access, to promote postoperative wound healing, or to minimize wound complications.

Suction-Assisted Lipectomy

Suction-assisted lipectomy of the abdominal area is a procedure in which excess fat deposits are removed from the trunk using a liposuction cannula with the goal of recontouring the body, thereby improving appearance. This procedure may be performed alone or as one component of the overall abdominoplasty or panniculectomy procedure. Suction-assisted lipectomy, when performed alone and not as part of a medically necessary abdominoplasty or panniculectomy, is considered cosmetic in nature. When the procedure is performed as part of a medically necessary abdominoplasty or panniculectomy, suction-assisted lipectomy of the trunk is considered incidental to the primary procedure.

Abdominal Wall Laxity and Back Pain

No correlation has been established between the presence of abdominal wall laxity or redundant pannus and the development of neck or back pain. There is insufficient evidence in the published, peer-reviewed scientific literature to support the use of abdominoplasty and/or panniculectomy to treat neck or back pain, including pain in the cervical, thoracic, lumbar or lumbosacral regions. Abdominoplasty or panniculectomy is considered not medically necessary when performed for the sole purpose of treating neck or back pain.

Professional Societies/Organizations

According to the ASPS practice parameter for abdominoplasty and panniculectomy, the procedures are most commonly performed for cosmetic indications. However, there are reconstructive indications such as abdominal wall defects, irregularities or pain caused by previous pelvic or lower abdominal surgery, umbilical hernias, intertriginous skin conditions and scarring (ASPS, 2007c). The ASPS recommended coverage criteria state that an abdominoplasty or panniculectomy should be considered a reconstructive procedure when performed to correct or relieve structural defects of the abdominal wall. When an abdominoplasty or panniculectomy is performed solely to enhance a patient's appearance in the absence of any signs or symptoms of functional abnormalities, the procedure should be considered cosmetic (ASPS, 2007a).

Summary

Abdominoplasty and panniculectomy are often performed together to achieve the best therapeutic result. These procedures are indicated for panniculitis that impairs function and is unresponsive to good personal hygiene and optimal medical management. Suction-assisted lipectomy and mini-abdominoplasty, when performed purely for cosmesis, are considered not medically necessary.

Coding/Billing Information

Note: This list of codes may not be all-inclusive.

Covered when medically necessary:

CPT®* Codes	Description
15830	Excision, excessive skin and subcutaneous tissue (includes lipectomy); abdomen, infraumbilical panniculectomy
15847	Excision, excessive skin and subcutaneous tissue (includes lipectomy); abdomen (eg, abdominoplasty) (includes umbilical transposition and fascial plication) (List separately in addition to code for primary procedure)

HCPCS Codes	Description
	No specific codes

ICD-9-CM Diagnosis Codes	Description
278.1	Localized adiposity
682.2	Other cellulitis and abscess, trunk
692.9	Contact dermatitis and other eczema, unspecified cause
695.89	Other specified erythematous conditions, other
701.8	Other specified hypertrophic and atrophic conditions of skin
701.9	Unspecified hypertrophic and atrophic conditions of skin
707.8	Chronic ulcer of other specified site
707.9	Chronic ulcer of unspecified site
729.30	Panniculitis, unspecified site
729.39	Panniculitis, unspecified; other site

Not Medically Necessary/Not Covered:

CPT* Codes	Description
158777 [†]	Suction assisted lipectomy; trunk

[†]**Note:** Not covered when performed alone and not as part of a medically necessary and covered abdominoplasty/panniculectomy procedure.

HCPCS Codes	Description
	No specific codes

ICD-9-CM Diagnosis Codes	Description
723.1	Cervicalgia
723.6	Panniculitis specified as affecting neck
724.2	Lumbago
724.5	Backache, unspecified
728.84	Diastasis of muscle
	Multiple/Varied

*Current Procedural Terminology (CPT®) © 2008 American Medical Association: Chicago, IL.

References

1. Aly A, Avila E, Cram AE. Endoscopic plastic surgery. *Surg Clin North Am*. 2000 Oct;80(5):1373-82.
2. American Society of Plastic Surgeons (ASPS). ASPS Recommended Insurance Coverage Criteria for Third-Party Payers. Abdominoplasty and Panniculectomy Unrelated to Obesity or Massive Weight Loss. 2006 Jul. Last updated 2007a Jan. Accessed Jan 10, 2007; Dec 29, 2007. Available at URL address: http://www.plasticsurgery.org/medical_professionals/health_policy/loader.cfm?url=/commonspot/security/getfile.cfm&PageID=18092
3. American Society of Plastic Surgeons (ASPS). ASPS Recommended Insurance Coverage Criteria for Third-Party Payers. Surgical Treatment of Skin Redundancy Following Massive Weight Loss. 2006 Jul. Last updated 2007b Jan. Accessed Jan 10, 2007; Dec 29, 2007. Available at URL address: http://www.plasticsurgery.org/medical_professionals/health_policy/loader.cfm?url=/commonspot/security/getfile.cfm&PageID=18091
4. American Society of Plastic Surgeons (ASPS). Practice Parameter for Abdominoplasty and Panniculectomy Unrelated to Obesity or Massive Weight Loss. 2006 Jul. Last updated 2007c Jan. Accessed Dec 29, 2007. Available at URL address: http://www.plasticsurgery.org/medical_professionals/health_policy/loader.cfm?url=/commonspot/security/getfile.cfm&PageID=18807
5. American Society of Plastic Surgeons (ASPS). Practice Parameter for Surgical Treatment of Skin Redundancy Following Massive Weight Loss. 2006 Jul. Last updated 2007d Jan. Accessed Dec 29, 2007. Available at URL address: http://www.plasticsurgery.org/medical_professionals/practice_manage/loader.cfm?url=/commonspot/security/getfile.cfm&PageID=18806
6. American Society of Plastic Surgeons (ASPS). Practice Advisory on Liposuction: Executive Summary. 2003 Mar. Accessed Jan 10, 2006; Jan 10, 2007; Dec 29, 2007. Available at URL address: http://www.plasticsurgery.org/medical_professionals/Policy_Statements/loader.cfm?url=/commonspot/security/getfile.cfm&PageID=7658
7. Arthurs ZM, Cuadrado D, Sohn V, Wolcott K, Lesperance K, Carter P, et al. Post-bariatric panniculectomy: pre-panniculectomy body mass index impacts the complication profile. *Am J Surg*. 2007 May;193(5):567-70; discussion 570.
8. Blomfield PI, Le T, Allen DG, Planner RS. Panniculectomy: a useful technique for the obese patient undergoing gynecological surgery. *Gynecol Oncol*. 1998 Jul;70(1):80-6.
9. Bragg TW, Jose RM, Srivastava S. Patient satisfaction following abdominoplasty: An NHS experience. *J Plast Reconstr Aesthet Surg*. 2007;60(1):75-8. Epub 2006 Sep 6.
10. Brauman D. Liposuction abdominoplasty: an evolving concept. *Plast Reconstr Surg*. 2003 Jul;112(1):288-98; discussion 299-301.
11. Chaouat M, Levan P, Lalanne B, Buisson T, Nicholau P, Mimoun M. Abdominal dermolipectomies: early postoperative complications and long-term unfavorable results. *Plast Reconstr Surg*. 2000 Dec;106(7):1614-8; discussion 1619-23.
12. Core GB, Mizgala CL, Bowen JC 3rd, Vasconez LO. Endoscopic abdominoplasty with repair of diastasis recti and abdominal wall hernia. *Clin Plast Surg*. 1995 Oct;22(4):707-22.
13. Dumanian GA, Denham W. Comparison of repair techniques for major incisional hernias. *Am J Surg*. 2003 Jan;185(1):61-5.
14. Fracalvieri M, Datta G, Bogetti P, Verna G, Pedrale R, Bocchiotti MA, et al. Abdominoplasty after weight loss in morbidly obese patients: a 4-year clinical experience. *Obes Surg*. 2007 Oct;17(10):1319-24.

15. Gmür RU, Banic A, Erni D. Is it safe to combine abdominoplasty with other dermolipectomy procedures to correct skin excess after weight loss? *Ann Plast Surg.* 2003 Oct;51(4):353-7.
16. Gravante G, Araco A, Sorge R, Araco F, Delogu D, Cervelli V. Wound infections in post-bariatric patients undergoing body contouring abdominoplasty: the role of smoking. *Obes Surg.* 2007 Oct;17(10):1325-31.
17. Hensel JM, Lehman JA Jr., Tantri MP, Parker MG, Wagner DS, Topham NS. An outcomes analysis and satisfaction survey of 199 consecutive abdominoplasties. *Ann Plast Surg.* 2001 Apr;46(4):357-63.
18. Hopkins MP, Shriner AM, Parker MG, Scott L. Panniculectomy at the time of gynecologic surgery in morbidly obese patients. *Am J Obstet Gynecol.* 2000 Jun;182(6):1502-5.
19. Iglesias M, Bravo L, Chavez-Munoz C, Barajas-Olivas A. Endoscopic abdominoplasty: an alternative approach. *Ann Plast Surg.* 2006 Nov;57(5):489-94.
20. Igwe D, Stanczyk M, Lee H, Felahy B, Tambi J, Fobi MA. Panniculectomy adjuvant to obesity surgery. *Obes Surg.* 2000 Dec;10(6):530-9.
21. Kaplan HY, Bar-Meir E. Safety of combining abdominoplasty and total abdominal hysterectomy: fifteen cases and review of the literature. *Ann Plast Surg.* 2005 Apr;54(4):390-2.
22. Lawrence N, Clark RE, Flynn TC, Coleman WP 3rd. American Society for Dermatologic Surgery guidelines of care for liposuction. *Dermatol Surg.* 2000 Mar;26(3):265-9.
23. Lockwood T. Rectus muscle diastasis in males: primary indication for endoscopically assisted abdominoplasty. *Plast Reconstr Surg.* 1998 May;101(6):1685-91; discussion 1692-4.
24. Logan JM, Broughton G 2nd. Plastic surgery: understanding abdominoplasty and liposuction. *AORN J.* 2008 Oct;88(4):587-600; quiz 601-4.
25. Manahan MA, Shermak MA. Massive panniculectomy after massive weight loss. *Plast Reconstr Surg.* 2006 Jun;117(7):2191-7; discussion 2198-9.
26. Mast BA. Safety and efficacy of outpatient full abdominoplasty. *Ann Plast Surg.* 2005 Mar;54(3):256-9.
27. Matarasso A. Liposuction as an adjunct to a full abdominoplasty revisited. *Plast Reconstr Surg.* 2000 Oct;106(5):1197-1202; discussion 1203-5.
28. Matarasso A, Belsley K. Abdominal contour procedures: evaluating the options. *Dermatol Clin.* 2005 Jul;23(3):475-93, vi-vii.
29. Nahas FX, Augusto SM, Ghelfond C. Should diastasis recti be corrected? *Aesthetic Plast Surg.* 1997 Jul-Aug;21(4):285-9.
30. Neaman KC, Hansen JE. Analysis of complications from abdominoplasty: a review of 206 cases at a university hospital. *Ann Plast Surg.* 2007 Mar;58(3):292-8.
31. Nguyen TT, Kim KA, Young RB. Tumescant mini abdominoplasty. *Ann Plast Surg.* 1997 Mar;38(3):209-12.
32. Powell JL, Kasperek DK, Connor GP. Panniculectomy to facilitate gynecologic surgery in morbidly obese women. *Obstet Gynecol.* 1999 Oct;94(4):528-31.
33. Ramirez OM. Abdominoplasty and abdominal wall rehabilitation: a comprehensive approach. *Plast Reconstr Surg.* 2000 Jan;105(1):425-35.

34. Ribeiro L, Accorsi AJ, Buss A. Midiabdominoplasty: indications and technique. *Aesthetic Plast Surg.* 1998 Sep-Oct;22(5):313-7.
35. Rogliani M, Silvi E, Labardi L, Maggiulli F, Cervelli V. Obese and nonobese patients: complications of abdominoplasty. *Ann Plast Surg.* 2006 Sep;57(3):336-8..
36. Shermak MA. Hernia repair and abdominoplasty in gastric bypass patients. *Plast Reconstr Surg.* 2006 Apr;117(4):1145-50; discussion 1151-2.
37. Sozer SO, Agullo FJ, Santillan AA, Wolf C. Decision Making in Abdominoplasty. *Aesthetic Plast Surg.* 2007 Jan 4; [Epub ahead of print]
38. Spiegelman JI, Levine RH. Abdominoplasty: a comparison of outpatient and inpatient procedures shows that it is a safe and effective procedure for outpatients in an office-based surgery clinic. *Plast Reconstr Surg.* 2006 Aug;118(2):517-22; discussion 523-4.
39. Stewart KJ, Stewart DA, Coghlan B, Harrison DH, Jones BM, Waterhouse N. Complications of 278 consecutive abdominoplasties. *J Plast Reconstr Aesthet Surg.* 2006;59(11):1152-5. Epub 2006 Jul 5.
40. Tillmanns TD, Kamelle SA, Abudayyeh I, McMeekin SD, Gold MA, Korkos TG, et al. Panniculectomy with simultaneous gynecologic oncology surgery. *Gynecol Oncol.* 2001 Dec;83(3):518-22.
41. Vila-Rovira R. Lipoabdominoplasty. *Clin Plast Surg.* 2008 Jan;35(1):95-104; discussion 105.
42. Yoho RA, Romaine JJ, O'Neil D. Review of the liposuction, abdominoplasty, and face-lift mortality and morbidity risk literature. *Dermatol Surg.* 2005 Jul;31(7 Pt 1):733-43; discussion 743.

Policy History

Pre-Merger Organizations	Last Review Date	Policy Number	Title
CIGNA HealthCare	2/15/2008	0027	Abdominoplasty and Panniculectomy
Great-West Healthcare	10/26/2006	95.287.06	Abdominoplasty and Panniculectomy
	7/19/2007	95.296.05	Diastasis Recti Repair

“CIGNA” and the “Tree of Life” logo are registered service marks of CIGNA Intellectual Property, Inc., licensed for use by CIGNA Corporation and its operating subsidiaries. All products and services are provided exclusively by such operating subsidiaries and not by CIGNA Corporation. Such operating subsidiaries include Connecticut General Life Insurance Company, CIGNA Behavioral Health, Inc., Intracorp, and HMO or service company subsidiaries of CIGNA Health Corporation and CIGNA Dental Health, Inc. In Arizona, HMO plans are offered by CIGNA HealthCare of Arizona, Inc. In California, HMO plans are offered by CIGNA HealthCare of California, Inc. and Great-West Healthcare of California, Inc. In Connecticut, HMO plans are offered by CIGNA HealthCare of Connecticut, Inc. In North Carolina, HMO plans are offered by CIGNA HealthCare of North Carolina, Inc. In Virginia, HMO plans are offered by CIGNA HealthCare Mid-Atlantic, Inc. All other medical plans in these states are insured or administered by Connecticut General Life Insurance Company.

Connecticut General Life Insurance Company has acquired the business of Great-West Healthcare from Great-West Life & Annuity Insurance Company (GWLA). Certain products continue to be provided by GWLA (Life, Accident and Disability, and Excess Loss). GWLA is not licensed to do business in New York. In New York, these products are sold by GWLA's subsidiary, First Great-West Life & Annuity Insurance Company, White Plains, N.Y.